

Nutrition and the Health of Young People

Centers for Disease Control and Prevention/Division of Adolescent and School Health

Diet and Disease

- Overweight and obesity, influenced by poor diet and inactivity, are significantly associated with an increased risk of diabetes, high blood pressure, high cholesterol, asthma, arthritis, and poor health status.¹
- Type 2 diabetes, formerly known as adult onset diabetes, has become increasingly prevalent among children and adolescents as rates of overweight and obesity rise.² A CDC study estimated that one in three American children born in 2000 will develop diabetes in their lifetime.³
- Overweight children and adolescents are more likely to become overweight or obese adults;⁴ one study showed that children who became overweight by age 8 were more severely obese as adults.⁵
- Early indicators of atherosclerosis, the most common cause of heart disease, begin as early as childhood and adolescence. Atherosclerosis is related to high blood cholesterol levels, which are associated with poor dietary habits.⁶
- Osteoporosis, a disease where bones become fragile and can break easily, is associated with inadequate intake of calcium.⁷

Overweight Among Youth

- Approximately 16% of children and adolescents ages 6-19 years are overweight.⁸
- The prevalence of overweight among children aged 6-11 years has more than doubled in the past 20 years and among adolescents aged 12-19 has more than tripled. Current estimates indicate that rates have not begun to decrease.^{8,9}
- While the prevalence of overweight and obesity has increased in all segments of the U.S. population, it is particularly common among minority groups.⁸

Prevalence of Overweight* Among Children and Adolescents in the United States, 1999-2002.⁹

Racial/Ethnic Group	Children (Ages 6 to 11)		Adolescents (Ages 12 to 19)	
	Male	Female	Male	Female
Black (Non-Hispanic)	17%	23%	19%	24%
Mexican American	27%	17%	25%	20%
White (Non-Hispanic)	14%	13%	15%	13%

Percent of Children and Adolescents Who Meet Dietary Guidelines, 1994-1996, 1998.¹⁰

Guideline	6-11 years old		12-19 years old	
	Boys	Girls	Boys	Girls
Fat	31%	34%	30%	35%
Saturated Fat	25%	25%	28%	34%
Vegetables	18%	19%	33%	26%
Fruits	23%	24%	14%	18%

* BMI at the 95th percentile or higher.



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION

March 2005



Diet and Academic Performance

- Research suggests that not having breakfast can affect children's intellectual performance.¹¹
- The percentage of young people who eat breakfast decreases with age; while 92% of children ages 6-11 eat breakfast, only 75-78% of adolescents ages 12-19 report eating breakfast.¹⁰

Eating Behaviors of Young People

- Less than 40% of children and adolescents in the United States meet the U.S. dietary guidelines for saturated fat.¹⁰
- Almost 80% of young people do not eat the recommended number of servings of fruits and vegetables.¹²
- Only 39% of children ages 2-17 meet the USDA's dietary recommendation for fiber (found primarily in dried beans and peas, fruits, vegetables, and whole grains).¹³
- Eighty-five percent of adolescent females do not consume enough calcium.⁷ During the last 25 years, consumption of milk, the largest source of calcium, has decreased 36% among adolescent females.¹⁴ Additionally, from 1978 to 1998, average daily soft drink consumption almost doubled among adolescent girls, increasing from 6 oz to 11 oz, and almost tripled among adolescent boys, from 7 oz to 19 oz.^{15,10}
- A large number of high school students use unsafe methods to lose or maintain weight. A nationwide survey found that during the 30 days preceding the survey 13% of students went without eating for one or more days; 6% had vomited or taken laxatives; and 9% had taken diet pills, powders, or liquids without the advice of their physicians.¹²

References

1. Mokdad AH, Ford ES, Bowman BA, Dietz WH, Vinicor F, Bales VS, Marks JS. Prevalence of obesity, diabetes, and obesity-related health risk factors, 2001. *Journal of the American Medical Association* 2003;289(1):76-79.
2. Rosenbloom AL, Joe JR, Young RS, Winter WE. Emerging epidemic of type 2 Diabetes in youth. *Diabetes Care* 1999;22(2):345-354.
3. Venkat Narayan KM, Boyle JP, Thompson TJ, Sorensen SW, Williamson DF. Lifetime risk for diabetes mellitus in the United States. *Journal of the American Medical Association* 2003;290(14):1884-1890.
4. Ferraro KF, Thorpe RJ Jr, Wilkinson JA. The life course of severe obesity: Does childhood overweight matter? *Journal of Gerontology* 2003;58B(2):S110-S119.
5. Freedman DS, Khan LK, Dietz WH, Srinivasan SR, Berenson GS. Relationship of childhood obesity to coronary heart disease risk factors in adulthood: the Bogalusa heart study. *Pediatrics* 2001;108(3):712-18.
6. Kavey RW, Daniels SR, Lauer RM, Atkins DL, Hayman LL, Taubert K. American Heart Association guidelines for primary prevention of atherosclerotic cardiovascular disease beginning in childhood. *Journal of Pediatrics* 2003;142(4):368-372.
7. National Institutes of Health. Preventing osteoporosis... why milk matters now for female teens. News and Features 1997:23. <http://www.nih.gov/news/nf/womenshealth/23.html>. (Accessed October 15, 2003)
8. Hedley AA, Ogden CL, Johnson CL, Carroll MD, Curtin LR, Flegal KM. Prevalence of overweight and obesity among US children, adolescents, and adults, 1999-2002. *Journal of the American Medical Association* 2004;291(23):2847-2850.
9. Ogden CL, Flegal KM, Carroll MD, Johnson CL. Prevalence and trends in overweight among US children and adolescents, 1999-2000. *Journal of the American Medical Association* 2002;288:1728-32.
10. United States Department of Agriculture. Continuing Survey of Food Intakes by Individuals 1994-96, 1998.
11. Pollitt E, Matthews R. Breakfast and cognition: an integrative summary. *American Journal of Clinical Nutrition* 1998;67(suppl):804S-813S.
12. Grunbaum JA, Kann L, Kinchen S, Ross J, Hawkins J, Lowry R, et al. Youth Risk Behavior Surveillance – United States, 2003. *Mortality and Morbidity Weekly Report* 2004;53(SS-2):1-95.
13. Lin BH, Guthrie J, Frazao E. American children's diets not making the grade. *Food Review* 2001;24(2):8-17.
14. Cavadini C, Siega-Riz AM, Popkin BM. US adolescent food intake trends from 1965 to 1996. *Archives of Disease in Childhood* 2000;83(1):18-24.
15. United States Department of Agriculture. Continuing Survey of Food Intakes by Individuals, 1987-88, Appendix A.

For Additional Information Contact:

CDC, Division of Adolescent and School Health
 4770 Buford Highway, NE
 Mail Stop K-12
 Atlanta, Georgia 30341-3717
 888-231-6405
 E-mail: HealthyYouth@cdc.gov